

Orthonormal Frame and $SO(3)$ Kaluza-Klein Dyon

Hsiang-Lin Lou^{*} and Pei-Hsuan Li[†]

Department of Physics, Tunghai University
Taichung, 407, Taiwan

November 9, 2009

Abstract

In previous paper, we present an $SO(3)$ Wu-Yang-like Kaluza-Klein dyon solution satisfies the Einstein equation in the seven-dimensional spacetimes. In this note, we will show an alternative approach using an orthonormal frame, the Cartan's structure equations, and calculating the affine spin connection one-form, curvature tensor and Ricci tensor. The results from these two different methods are coincident.

PACS numbers: 04.50.Cd

^{*} Email: slou@thu.edu.tw

[†] Email: ludmila.phys@gmail.com

1 Introduction

In previous paper,[1] we have shown that a new $SO(3)$ Wu-Yang-like [2] seven-dimensional Kaluza-Klein(KK) dyon solution satisfies the Einstein equation, by calculating the Christoffel symbols and the Ricci tensor. In this note, we will show an alternative approach [3] using an orthonormal frame, the Cartan's structure equations and calculating the affine spin connection one-form, curvature tensor and Ricci tensor.

Now consider the Kaluza-Klein theory [4] in a $(4 + N)$ -manifold M with a metric $\bar{g}_{AB}(x)$ on M in local coordinates $\bar{x}^A$. The line element is

$$d\bar{s}^2 = \bar{g}_{AB}d\bar{x}^A d\bar{x}^B \quad (1)$$

$$= g_{\mu\nu}(x)dx^\mu dx^\nu + \gamma_{mn}(y)(dy^m + B_\mu^m dx^\mu)(dy^n + B_\nu^n dx^\nu), \quad (2)$$

where x parametrizes four-dimensional spacetimes, y parametrizes extra dimensions. We use $A, B, C \dots$ indices to represent the total spacetimes; $\mu, \nu, \rho \dots$ to represent the four-dimensional spacetimes; $m, n, l \dots$ to represent the extra dimensions. $g_{\mu\nu}$ is only a function of x , and γ_{mn} is only a function of y .

The ansatz [1] of the Kaluza-Klein dyonic metric admitting $SO(3)$ Killing vectors is

$$\begin{aligned} d\bar{s}^2 = & - e^{2\Psi} dt^2 \\ & + e^{2\Lambda} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \\ & + (dR + B_\mu^5 dx^\mu)^2 + R^2 (d\Theta + B_\mu^6 dx^\mu)^2 + R^2 \sin^2 \Theta (d\Phi + B_\mu^7 dx^\mu)^2. \end{aligned} \quad (3)$$

r, θ, ϕ are three coordinates of the ordinary three-dimensional spherical coordinate system, $(r, \theta, \phi) = (\bar{x}^1, \bar{x}^2, \bar{x}^3) = (x^1, x^2, x^3)$. R, Θ, Φ are another three coordinates of the spherical coordinate system in the extra dimensions, $(R, \Theta, \Phi) = (\bar{x}^5, \bar{x}^6, \bar{x}^7) = (y^5, y^6, y^7)$. Ψ and Λ are two functions of r . We have $g_{00} = -e^{2\Psi}$, $g_{11} = e^{2\Lambda}$, $g_{22} = r^2$, $g_{33} = r^2 \sin^2 \theta$, $\gamma_{55} = 1$, $\gamma_{66} = R^2$, $\gamma_{77} = R^2 \sin^2 \Theta$.

B_μ^m cannot be identified as the Yang-Mills field. To extract the true Yang-Mills field, one has to introduce the Killing vectors

$$L_a \equiv -i\zeta_a^m \partial_m, \quad (4)$$

which generate an $SO(3)$ algebra,

$$[L_a, L_b] = i\epsilon_{abc} L_c, \quad (5)$$

Inserting L_a of (4) into equation (5), one gets the Killing's equation

$$\zeta_a^m \partial_m \zeta_b^n - \zeta_b^m \partial_m \zeta_a^n = -\epsilon_{abc} \zeta_c^n. \quad (6)$$

With these Killing vectors, one can define

$$B_\mu^m = \zeta_a^m A_\mu^a, \quad (7)$$

where A_μ^a is the true Yang-Mills field and ζ_a^m is only a function of y . Defining

$$\tilde{\mathcal{F}}_{\mu\nu}^m \equiv \partial_\mu B_\nu^m - B_\nu^l \partial_l B_\mu^m - (\mu \leftrightarrow \nu) \quad (8)$$

$$= \zeta_a^m F_{\mu\nu}^a, \quad (9)$$

where $F_{\mu\nu}^a$ is the true field strength tensor of the Yang-Mills field,

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + \epsilon_{abc} A_\mu^b A_\nu^c. \quad (10)$$

It can be checked that the components of ζ_a^m ,

$$\zeta_a^5 = 0, \quad (11)$$

$$\zeta_a^6 = \hat{\Phi}_a, \quad (12)$$

$$\zeta_a^7 = -\frac{1}{\sin \Theta} \hat{\Theta}_a, \quad (13)$$

satisfy the Killing equation (6). The gauge-field components of the Wu-Yang-like KK dyon are

$$A_0^a = \frac{1}{r} \hat{r}^a, \quad A_1^a = 0, \quad (14)$$

$$A_2^a = -\hat{\phi}^a, \quad A_3^a = \sin \theta \hat{\theta}^a. \quad (15)$$

A_1^a, A_2^a, A_3^a are just the spherical coordinate representation of the Wu-Yang monopole field in the ordinary gauge theory of four-dimensional spacetimes. The electric field of the KK dyon is

$$F_{01}^a = \frac{1}{r^2} \hat{r}^a, \quad (16)$$

while the magnetic field is

$$F_{23}^a = -\sin \theta \hat{r}^a. \quad (17)$$

The fields B_μ^m in (7) can be rewritten as

$$B_\mu^5 = 0, \quad B_1^m = 0, \quad (18)$$

$$B_0^6 = \frac{1}{r} \hat{r} \cdot \hat{\Phi}, \quad B_0^7 = -\frac{1}{r \sin \Theta} \hat{r} \cdot \hat{\Theta}, \quad (19)$$

$$B_2^6 = -\hat{\phi} \cdot \hat{\Phi}, \quad B_2^7 = \frac{1}{\sin \Theta} \hat{\phi} \cdot \hat{\Theta}, \quad (20)$$

$$B_3^6 = \sin \theta \hat{\theta} \cdot \hat{\Phi}, \quad B_3^7 = -\frac{\sin \theta}{\sin \Theta} \hat{\theta} \cdot \hat{\Theta}. \quad (21)$$

The nonzero components of $\tilde{\mathcal{F}}_{\mu\nu}^m$ are

$$\tilde{\mathcal{F}}_{01}^6 = \frac{1}{r^2} \hat{r} \cdot \hat{\Phi}, \quad \tilde{\mathcal{F}}_{01}^7 = \frac{-1}{r^2 \sin \Theta} \hat{r} \cdot \hat{\Theta}, \quad (22)$$

$$\tilde{\mathcal{F}}_{23}^6 = -\sin \theta \hat{r} \cdot \hat{\Phi}, \quad \tilde{\mathcal{F}}_{23}^7 = \frac{\sin \theta}{\sin \Theta} \hat{r} \cdot \hat{\Theta}. \quad (23)$$

2 Orthonormal Frame

We now decompose the metric into vielbeins as

$$\bar{g}_{AB} = \eta_{\bar{a}\bar{b}} e^{\bar{a}}_A e^{\bar{b}}_B, \quad \bar{a}, \bar{b} = 0, 1, 2, 3, 5, 6, 7, \quad (24)$$

where the $\eta_{\bar{a}\bar{b}}$ is a seven-dimensional flat Minkowski space metric,

$$\eta_{\bar{a}\bar{b}} = \text{diag}(-1, 1, 1, 1, 1, 1, 1) \quad (25)$$

The inverse of $e^{\bar{a}}_A$ is defined by

$$E^{\bar{a}A} = \eta_{\bar{a}\bar{b}} \bar{g}^{AB} e^{\bar{b}}_B \quad (26)$$

which obeys

$$E^{\bar{a}A} e^{\bar{b}}_A = \delta^{\bar{b}}_{\bar{a}}, \quad \eta^{\bar{a}\bar{b}} E^{\bar{a}A} E^{\bar{b}B} = \bar{g}^{AB}. \quad (27)$$

$e^{\bar{a}}_A$ is the matrix which transforms the coordinate basis $d\bar{x}^A$ of the cotangent space $T_x^*(M)$ to an orthonormal basis of the same space $T_x^*(M)$,

$$e^{\bar{a}} = e^{\bar{a}}_A d\bar{x}^A. \quad (28)$$

The vielbein basis of the SO(3) KK dyonic metric can be written as

$$e^0 = e^\Psi dt, \quad e^1 = e^\Lambda dr, \quad e^2 = r d\theta, \quad e^3 = r \sin\theta d\phi, \quad (29)$$

$$e^5 = dR, \quad e^6 = R(d\Theta + B_\mu^6 dx^\mu), \quad e^7 = R \sin\Theta (d\Phi + B_\mu^7 dx^\mu). \quad (30)$$

The affine spin connection one-form $\omega^{\bar{a}}_{\bar{b}}$ are introduced by

$$de^{\bar{a}} + \omega^{\bar{a}}_{\bar{b}} \wedge e^{\bar{b}} = 0 \quad (31)$$

and the metricity condition

$$\omega_{\bar{a}\bar{b}} = -\omega_{\bar{b}\bar{a}}. \quad (32)$$

The curvature 2-form is defined as

$$R^{\bar{a}}_{\bar{b}} = d\omega^{\bar{a}}_{\bar{b}} + \omega^{\bar{a}}_{\bar{c}} \wedge \omega^{\bar{c}}_{\bar{b}} = \bar{R}^{\bar{a}}_{\bar{b}\bar{c}\bar{d}} e^{\bar{c}} \wedge e^{\bar{d}}. \quad (33)$$

Equations (31) and (33) are called Cartan's structure equations. The components of the curvature tensor have the relations,

$$\bar{R}_{\bar{a}\bar{b}\bar{c}\bar{d}} = -\bar{R}_{\bar{b}\bar{a}\bar{c}\bar{d}} = -\bar{R}_{\bar{a}\bar{b}\bar{d}\bar{c}} = \bar{R}_{\bar{c}\bar{d}\bar{a}\bar{b}}. \quad (34)$$

and satisfy the Bianchi identity,

$$\bar{R}_{\bar{a}\bar{b}\bar{c}\bar{d}} + \bar{R}_{\bar{a}\bar{c}\bar{d}\bar{b}} + \bar{R}_{\bar{a}\bar{d}\bar{b}\bar{c}} = 0. \quad (35)$$

The components of the affine spin connection one-form $\omega^{\bar{a}}_{\bar{b}}$ of the $SO(3)$ KK dyon can be written more explicitly as

$$\omega^0_1 = \Psi' e^{-\Lambda} e^0 + \frac{R}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Phi}) e^6 - \frac{R}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Theta}) e^7, \quad (36)$$

$$\omega^0_2 = 0, \quad \omega^0_3 = 0, \quad \omega^0_5 = 0, \quad (37)$$

$$\omega^0_6 = \frac{R}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Phi}) e^1, \quad (38)$$

$$\omega^0_7 = -\frac{R}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Theta}) e^1, \quad (39)$$

$$\omega^1_2 = -\frac{1}{r} e^{-\Lambda} e^2, \quad (40)$$

$$\omega^1_3 = -\frac{1}{r} e^{-\Lambda} e^3, \quad \omega^1_5 = 0, \quad (41)$$

$$\omega^1_6 = \frac{R}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Phi}) e^0 \quad (42)$$

$$\omega^1_7 = -\frac{R}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Theta}) e^0 \quad (43)$$

$$\omega^2_3 = -\frac{1}{r} \cot\theta e^3 + \frac{R}{2r^2} (\hat{r} \cdot \hat{\Phi}) e^6 - \frac{R}{2r^2} (\hat{r} \cdot \hat{\Theta}) e^7, \quad \omega^2_5 = 0, \quad (44)$$

$$\omega^2_6 = \frac{R}{2r^2} (\hat{r} \cdot \hat{\Phi}) e^3, \quad (45)$$

$$\omega^2_7 = -\frac{R}{2r^2} (\hat{r} \cdot \hat{\Theta}) e^3, \quad \omega^3_5 = 0, \quad (46)$$

$$\omega^3_6 = -\frac{R}{2r^2} (\hat{r} \cdot \hat{\Phi}) e^2, \quad \omega^3_7 = \frac{R}{2r^2} (\hat{r} \cdot \hat{\Theta}) e^2, \quad (47)$$

$$\omega^5_6 = -\frac{1}{R} e^6, \quad \omega^5_7 = -\frac{1}{R} e^7, \quad (48)$$

$$\begin{aligned} \omega^6_7 = & -\frac{1}{R} \cot\theta e^7 - \frac{1}{r} e^{-\Psi} \cot\theta (\hat{r} \cdot \hat{\Theta}) e^0 - \frac{1}{r} e^{-\Psi} (\hat{r} \cdot \hat{R}) e^0 \\ & + \frac{1}{r} \cot\theta (\hat{\Phi} \cdot \hat{\Theta}) e^2 + \frac{1}{r} (\hat{\Phi} \cdot \hat{R}) e^2 - \frac{1}{r} \cot\theta (\hat{\theta} \cdot \hat{\Theta}) e^3 - \frac{1}{r} (\hat{\theta} \cdot \hat{R}) e^3. \end{aligned} \quad (49)$$

The nonzero components of the curvature tensor are

$$\bar{R}_{0101} = e^{-2\Lambda} (\Psi'' - \Psi' \Lambda' + (\Psi')^2) - \frac{3}{4} \frac{R^2}{r^4} e^{-2\Psi-2\Lambda} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}, \quad (50)$$

$$\bar{R}_{0116} = \frac{R}{2r^2} e^{-\Psi-2\Lambda} (\Psi' + \Lambda' + \frac{2}{r}) (\hat{r} \cdot \hat{\Phi}), \quad (51)$$

$$\bar{R}_{0117} = -\frac{R}{2r^2} e^{-\Psi-2\Lambda} (\Psi' + \Lambda' + \frac{2}{r}) (\hat{r} \cdot \hat{\Theta}), \quad (52)$$

$$\bar{R}_{0123} = \frac{R^2}{2r^4} e^{-\Psi-\Lambda} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}, \quad (53)$$

$$\bar{R}_{0156} = -\frac{1}{r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{0157} = \frac{1}{r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Theta}), \quad (54)$$

$$(55)$$

$$\bar{R}_{0167} = -\frac{1}{r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{R}), \quad \bar{R}_{0202} = \frac{1}{r} \Psi' e^{-2\Lambda}, \quad (56)$$

$$\bar{R}_{0213} = \frac{R^2}{4r^4} e^{-\Psi-\Lambda} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}, \quad (57)$$

$$\bar{R}_{0226} = -\frac{R}{2r^3} e^{-\Psi-2\Lambda} (\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{0227} = \frac{R}{2r^3} e^{-\Psi-2\Lambda} (\hat{r} \cdot \hat{\Theta}), \quad (58)$$

$$\bar{R}_{0303} = \frac{1}{r} \Psi' e^{-2\Lambda}, \quad (59)$$

$$\bar{R}_{0312} = -\frac{R^2}{4r^4} e^{-\Psi-\Lambda} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}, \quad (60)$$

$$\bar{R}_{0336} = -\frac{R}{2r^3} e^{-\Psi-2\Lambda} (\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{0337} = \frac{R}{2r^3} e^{-\Psi-2\Lambda} (\hat{r} \cdot \hat{\Theta}), \quad (61)$$

$$\bar{R}_{0516} = -\frac{1}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{0517} = \frac{1}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Theta}), \quad (62)$$

$$\bar{R}_{0606} = \frac{R^2}{4r^4} e^{-2\Psi-2\Lambda} (\hat{r} \cdot \hat{\Phi})^2, \quad \bar{R}_{0607} = -\frac{R^2}{4r^4} e^{-2\Psi-2\Lambda} (\hat{r} \cdot \hat{\Theta})(\hat{r} \cdot \hat{\Phi}), \quad (63)$$

$$\bar{R}_{0615} = \frac{1}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{0617} = -\frac{1}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{R}), \quad (64)$$

$$\bar{R}_{0707} = \frac{R^2}{4r^4} e^{-2\Psi-2\Lambda} (\hat{r} \cdot \hat{\Theta})^2, \quad \bar{R}_{0715} = -\frac{1}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{\Theta}), \quad (65)$$

$$\bar{R}_{0716} = \frac{1}{2r^2} e^{-\Psi-\Lambda} (\hat{r} \cdot \hat{R}), \quad \bar{R}_{1212} = \frac{1}{r} \Lambda' e^{-2\Lambda}, \quad (66)$$

$$\bar{R}_{1236} = \frac{R}{2r^3} e^{-\Lambda} (\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{1237} = -\frac{R}{2r^3} e^{-\Lambda} (\hat{r} \cdot \hat{\Theta}), \quad (67)$$

$$\bar{R}_{1313} = \frac{1}{r} \Lambda' e^{-2\Lambda}, \quad \bar{R}_{1326} = -\frac{R}{2r^3} e^{-\Lambda} (\hat{r} \cdot \hat{\Phi}), \quad (68)$$

$$\bar{R}_{1327} = \frac{R}{2r^3} e^{-\Lambda} (\hat{r} \cdot \hat{\Theta}), \quad \bar{R}_{1616} = -\frac{R^2}{4r^4} e^{-2\Psi-2\Lambda} (\hat{r} \cdot \hat{\Phi})^2, \quad (69)$$

$$\bar{R}_{1617} = \frac{R^2}{4r^4} e^{-2\Psi-2\Lambda} (\hat{r} \cdot \hat{\Theta})(\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{1623} = -\frac{R}{r^3} e^{-\Lambda} (\hat{r} \cdot \hat{\Phi}), \quad (70)$$

$$\bar{R}_{1717} = -\frac{R^2}{4r^4} e^{-2\Psi-2\Lambda} (\hat{r} \cdot \hat{\Theta})^2, \quad \bar{R}_{1723} = \frac{R}{r^3} e^{-\Lambda} (\hat{r} \cdot \hat{\Theta}), \quad (71)$$

$$\bar{R}_{2323} = \frac{1}{r^2} (1 - e^{-2\Lambda}) - \frac{3}{4} \frac{R^2}{r^4} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}, \quad (72)$$

$$\bar{R}_{2356} = \frac{1}{r^2} (\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{2367} = \frac{1}{r^2} (\hat{r} \cdot \hat{R}), \quad (73)$$

$$\bar{R}_{2357} = -\frac{1}{r^2} (\hat{r} \cdot \hat{\Theta}), \quad \bar{R}_{2536} = \frac{1}{2r^2} (\hat{r} \cdot \hat{\Phi}), \quad (74)$$

$$\bar{R}_{2537} = -\frac{1}{2r^2} (\hat{r} \cdot \hat{\Theta}), \quad \bar{R}_{2626} = \frac{R^2}{4r^4} (\hat{r} \cdot \hat{\Phi})^2 \quad (75)$$

$$\bar{R}_{2627} = -\frac{R^2}{4r^4} (\hat{r} \cdot \hat{\Theta})(\hat{r} \cdot \hat{\Phi}), \quad \bar{R}_{2635} = -\frac{1}{2r^2} (\hat{r} \cdot \hat{\Phi}) \quad (76)$$

$$\bar{R}_{2637} = \frac{1}{2r^2} (\hat{r} \cdot \hat{R}), \quad \bar{R}_{2727} = \frac{R^2}{4r^4} (\hat{r} \cdot \hat{\Theta})^2 \quad (77)$$

$$\bar{R}_{2735} = \frac{1}{2r^2} (\hat{r} \cdot \hat{\Theta}), \quad \bar{R}_{2736} = -\frac{1}{2r^2} (\hat{r} \cdot \hat{R}) \quad (78)$$

$$\bar{R}_{3636} = \frac{R^2}{4r^4} (\hat{r} \cdot \hat{\Phi})^2, \quad \bar{R}_{3637} = -\frac{R^2}{4r^4} (\hat{r} \cdot \hat{\Theta})(\hat{r} \cdot \hat{\Phi}), \quad (79)$$

$$\bar{R}_{3737} = \frac{R^2}{4r^4} (\hat{r} \cdot \hat{\Theta})^2. \quad (80)$$

We have tacitly omitted nearly one half of the nonzero components of the curvature tensor, because of the relations in (34), for shorthand.

3 Einstein Equation

Contracting the curvature tensor, one will get the Ricci tensor

$$\bar{R}_{\bar{a}\bar{b}} = \eta^{\bar{c}\bar{d}} \bar{R}_{\bar{c}\bar{a}\bar{d}\bar{b}}. \quad (81)$$

Substituting the components of the dyonic metric,

$$e^{2\Psi} = 1 - \frac{r_s}{r}, \quad (82)$$

$$e^{2\Lambda} = \left(1 - \frac{r_s}{r}\right)^{-1}, \quad (83)$$

where r_s is the Schwarzschild radius, then we can obtain almost $\bar{R}_{\bar{a}\bar{b}}$ are zero except

$$\bar{R}_{00} = -\frac{1}{2} \frac{R^2}{r^4} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}, \quad (84)$$

$$\bar{R}_{11} = +\frac{1}{2} \frac{R^2}{r^4} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}, \quad (85)$$

$$\bar{R}_{22} = -\frac{1}{2} \frac{R^2}{r^4} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}, \quad (86)$$

$$\bar{R}_{33} = -\frac{1}{2} \frac{R^2}{r^4} \{(\hat{r} \cdot \hat{\Theta})^2 + (\hat{r} \cdot \hat{\Phi})^2\}. \quad (87)$$

Then one has the Ricci scalar curvature,

$$\bar{R} = \bar{\eta}^{\bar{a}\bar{b}} \bar{R}_{\bar{a}\bar{b}} \quad (88)$$

$$= -\bar{R}_{00} + \bar{R}_{11} + \bar{R}_{22} + \bar{R}_{33} \quad (89)$$

$$= 0. \quad (90)$$

From the fields $\tilde{\mathcal{F}}_{\mu\nu}^m$ in (22) and (23), the components of the Ricci tensor, (84) ~ (87), can be recast into the form,

$$\bar{R}_{\bar{a}\bar{b}} = -\frac{1}{2} E_{\bar{a}}^{\mu} E_{\bar{b}}^{\nu} \bar{g}^{\alpha\beta} \gamma_{mn} \tilde{\mathcal{F}}_{\mu\alpha}^m \tilde{\mathcal{F}}_{\nu\beta}^n, \quad (91)$$

or

$$\bar{R}_{\mu\nu} = -\frac{1}{2}\bar{g}^{\alpha\beta}\gamma_{mn}\tilde{\mathcal{F}}_{\mu\alpha}^m\tilde{\mathcal{F}}_{\nu\beta}^n. \quad (92)$$

Since the identity, $\gamma_{mn}\tilde{\mathcal{F}}_{\mu\nu}^m\tilde{\mathcal{F}}^{\mu\nu n} = 0$, holds, the right-hand side of the equation (92) can be identified as 8π times the stress-energy tensor of the Yang-Mills field,

$$\bar{R}_{\mu\nu} = 8\pi\bar{T}_{\mu\nu}, \quad \bar{T}_{\mu\nu} = \frac{-1}{16\pi}\bar{g}^{\alpha\beta}\gamma_{mn}\tilde{\mathcal{F}}_{\mu\alpha}^m\tilde{\mathcal{F}}_{\nu\beta}^n. \quad (93)$$

Then the Einstein equation, $\bar{R}_{AB} - \frac{1}{2}\bar{g}_{AB}\bar{R} = 8\pi\bar{T}_{AB}$, is satisfied, where some components of $\bar{T}_{AB}$ are zero, $\bar{T}_{\mu m} = 0$ and $\bar{T}_{mn} = 0$.

The $SO(3)$ KK dyonic metric satisfies the Einstein equation in the seven-dimensional. The results from two different methods are coincident. .

References

- [1] Hsiang-Lin Lou and Pei-Hsuan Li, *Nonabelian Kaluza-Klein Dyon*, arXiv:0910.2023
- [2] T.T. Wu and C.N. Yang, in " *Properties of Matter under Unusual Conditions*" edited by H. Mark and Fernbach, p.349, (1968) (New York, Interscience)
- [3] T.Eguchi, P.B.Gilkey and A.J.Hanson *Gravitation, Gauge Theories and Differential Geometry*, Phys. Rep.**66**, (1980) 213.
- [4] J.M.Overduin and P.S.Wesson, *Kaluza-Klein Gravity*, Phys. Rep. **283**, (1997) 303.